

Exercise 1 for the lecture

NUMERICS III

SoSe 2018

http://numerik.mi.fu-berlin.de/wiki/SS_2018/NumericsIII.php

Due: Wed, 02-05-2018

Problem 1 (4 TP)

Consider the space $V = \{f \in C(\mathbb{R}) \mid \|f\|_\infty < \infty\}$ with norm $\|\cdot\|_\infty$. Let $u_n(x) = \max\{0, 1 - |xn|\}$ for all $n \in \mathbb{N}$ and $x \in \mathbb{R}$. Show that $(u_n)_{n \in \mathbb{N}}$ has no converging subsequence in V and conclude that the unit sphere in V is not compact.

Problem 2 (4 TP)

Show that the Laplace operator is invariant under translations and rotations of the coordinate system: Let $A \in \mathbb{R}^{d \times d}$ be an orthogonal matrix with columns $a_i \in \mathbb{R}^d$. Given a rotation around a center $c \in \mathbb{R}^d$, defined by $\psi(x) = A(x - c) + c$, prove that the sequence of identities

$$\Delta u(x) = (\Delta(u \circ \psi))(\psi^{-1}x) = \Delta_A u(x) := \sum_{i=1}^d \frac{\partial^2}{\partial a_i \partial a_i} u(x)$$

holds for all $x \in \mathbb{R}^d$.

Problem 3 (5 TP)

For all $\varepsilon \in \mathbb{R}_{>0}$ let

$$f_\varepsilon(x) = \begin{cases} 1 & , x \in [-1, -\varepsilon] \\ -\frac{x}{\varepsilon} & , x \in (-\varepsilon, \varepsilon) \\ -1 & , x \in [\varepsilon, 1] \end{cases} \quad u_\varepsilon(x) = \begin{cases} -\frac{1}{2}x^2 - \frac{\varepsilon^2+3}{6}x - \frac{\varepsilon^2}{6} & , x \in [-1, -\varepsilon] \\ \frac{1}{6\varepsilon}x^3 + \frac{3\varepsilon-\varepsilon^2-3}{6}x & , x \in (-\varepsilon, \varepsilon) \\ \frac{1}{2}x^2 - \frac{\varepsilon^2+3}{6}x + \frac{\varepsilon^2}{6} & , x \in [\varepsilon, 1]. \end{cases}$$

a) Verify that u_ε solves

$$-\Delta u_\varepsilon = f_\varepsilon \text{ on } (-1, 1), \quad u_\varepsilon = 0 \text{ on } \{-1, 1\}$$

in the classical sense.

b) Prove that u_ε converges pointwise to

$$u(x) = \begin{cases} -\frac{x}{2}(x+1) & , x \in [-1, 0] \\ \frac{x}{2}(x-1) & , x \in [0, 1], \end{cases}$$

for $\varepsilon \rightarrow 0$.

c) Show that u_ε does not converge to u in $C^2([-1, 1])$.

d) Let $\|v\|_{H^1} := \sqrt{\int_{-1}^1 (v'(x))^2 dx}$. Show that $\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u\|_{H^1} = 0$.

e) Let

$$f(x) = \begin{cases} 1 & \text{for } x \leq 0 \\ -1 & \text{for } x > 0. \end{cases}$$

Prove that u is a weak solution of

$$-\Delta u = f \text{ on } (-1, 1), \quad u = 0 \text{ on } \{-1, 1\}$$

by verifying the variational equality

$$\forall v \in C_0^\infty((-1, 1)): \int_{-1}^1 u'(x) v'(x) dx = \int_{-1}^1 f(x) v(x) dx.$$

GENERAL REMARKS

You have to do the exercises in groups of up 3 people. Be prepared to demonstrate your solutions to theoretical problems at the given date in the tutorial. Solutions for programming problems have to be submitted via e-mail to adjurdjevac@mi.fu-berlin.de with a subject starting by [NumericsII] and denoting all members of the group. Please follow the additional advise for programming exercises on the homepage.