

Exercise 8 for the lecture  
NUMERICS II  
WS 2011/12

**Due: till Thursday, 5. January 12 o'clock**

**Problem 1** (4 TP)

The pendulum of length  $L$  and mass  $m$  is a simple mechanical system. We want to consider it in the Cartesian coordinates. It's behaviour is described by the below DAE containing two ODEs of second order and a position constraint

$$\begin{aligned}x'' &= -\lambda x \\y'' &= -\lambda y - g \\0 &= x^2 + y^2 - L^2\end{aligned}$$

where  $g$  is the constant gravity,  $\lambda$  the Lagrange multiplier and is unknown.

- Rewrite these two ODEs of the second order into four ODEs of the first order.
- Compute the differentiation index of this pendulum problem.

**Problem 2** (3 TP)

Consider the initial value problem

$$\begin{aligned}y'(t) &= f(y(t), z(t)), & y(0) &= y_0, \\ \epsilon z'(t) &= g(y(t), z(t)), & z(0) &= z_0\end{aligned}$$

with

$$f(y, z) = yz, \quad g(y, z) = z - z^3.$$

For which solutions  $y^*, z^*$  of  $g(y^*, z^*) = 0$  does

$$\max_{\lambda} \operatorname{Re}(\lambda) < 0, \quad \lambda \in \sigma \left( \frac{dg}{dz}(y^*, z^*) \right)$$

hold? Solve the corresponding DAE system for  $\epsilon = 0$  with consistent initial values. Discuss the asymptotic behavior of the singularly perturbed problem with  $\epsilon > 0$  for  $t \rightarrow \infty$ .

**Problem 3** (3 TP)

Consider the DAE

$$Nz'(t) = z(t) + f(t) \quad (1)$$

with  $f \in C^\infty(\mathbb{R}, \mathbb{R}^d)$ . Show that if  $N$  is nilpotent of degree  $\nu$  then  $\nu$ -fold derivation of (1) leads to an ODE without algebraic constraints. What is the differentiation index of (1) in this case ?

**Problem 4** (2 TP + 4 PP)

Consider the Schmitt-trigger initial value problem

$$\begin{pmatrix} C_J & 0 & -C_J & 0 & 0 \\ 0 & C_0 & 0 & -C_0 & 0 \\ -C_J & 0 & 2C_J & -C_J & 0 \\ 0 & -C_0 & -C_J & C_0 + C_J & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} u' = - \begin{pmatrix} G_1 u_1 + (1 - \alpha)g(u_1 - u_3) \\ G_2 u_2 + G_4(u_2 - u_4) + \alpha g(u_1 - u_3) \\ G_3 u_3 - g(u_1 - u_3) - g(u_4 - u_3) \\ G_4(u_4 - u_2) + (1 - \alpha)g(u_4 - u_3) \\ G_5 u_5 + \alpha g(u_4 - u_3) \end{pmatrix} + \begin{pmatrix} G_1 V_{in} \\ G_2 V_{DD} \\ 0 \\ 0 \\ G_5 V_{DD} \end{pmatrix}.$$

- a) Transform this problem to the semi-explicit form.
- b) Find a consistent initial value for the parameters

$$G = (200, 1600, 100, 3200, 1600), \quad C_J = 10^{-12}, \quad C_0 = 40 \cdot 10^{-12},$$

$$g(x) = 10^{-6} \left( \exp\left(\frac{x}{0.026}\right) - 1 \right), \quad \alpha = 0.99, \quad V_{dd} = 1$$

and the input function

$$V_{in}(t) = 2 \sin(2\pi t) + 0.2 \sin(20\pi t).$$

- c) Solve the problem with the above parameters and initial value numerically on the interval  $[0, 2]$  using an appropriate MATLAB method.
- d) Solve the problem with the above parameters and initial value numerically on the interval  $[0, 2]$  with the state-space method using the Runge-Kutta-4 method.

(Mind the minus sign in the right hand side !)