

Exercise 9 for the lecture
NUMERICS II
WS 2011/12

Due: till Thursday, 12. January 12 o'clock

Problem 1 (5 TP)

- a) Show that the symplectic Euler method is symplectic.
- b) Show that the trapezoidal rule is symplectic if the Hamiltonian is quadratic, i.e. $H(y) = y^T C y$ holds with a symmetric real matrix C .
- c) Show that the trapezoidal rule is not symplectic in general.

Problem 2 (4 TP)

Consider the system $q' = p; p' = f(q)$.
The explicit one-step method given by

$$\begin{aligned}p_{n+\frac{1}{2}} &= p_n + \frac{\tau}{2} f(q_n) \\q_{n+1} &= q_n + \tau p_{n+\frac{1}{2}} \\p_{n+1} &= p_{n+\frac{1}{2}} + \frac{\tau}{2} f(q_{n+1})\end{aligned}$$

is called Störmer-Verlet method.

Show that the Störmer-Verlet method is symplectic and has second order.

Problem 3 (5 PP)

Consider the pendulum equation in polar coordinates

$$\begin{pmatrix} q' \\ p' \end{pmatrix} = \begin{pmatrix} \frac{1}{m} p \\ -m \frac{g}{r_0} \cos q \end{pmatrix} \quad \begin{pmatrix} q \\ p \end{pmatrix} (0) = \begin{pmatrix} q_0 \\ p_0 \end{pmatrix}$$

where q , g , m , and r_0 denote the angle, the gravity, the mass and the radius, respectively.

- a) Implement the Störmer-Verlet method for this equation in `matlab` as function `[p, q, t] = StörmerVerlet(m, g, r0, p0, q0, I, tau)`, where (m, g, r_0) , (p_0, q_0) , I , and τ denote the problem parameters, the initial values, the time interval and the step size, respectively.
- b) Test your program with the radius $r_0 = 10\text{cm}$, the mass $m = 100\text{g}$, and the gravity of the moon. Use the time interval $[0\text{s}, 20\text{s}]$ with the initial value $p_0 = 0 \frac{\text{kgm}}{\text{s}}$, $q_0 = 0\text{m}$ for various time step sizes.
- c) Plot the solution in the phase space, the solution in Euclidean coordinates, and the associated Hamiltonian H .